

Primary Cells for Scientific Research

Mouse Hepatic Stellate Cells (Primary cell immortalization)

Cat. No. : CP-M041Y

General Information

Species	Mouse/C57BL/6 mouse
Tissue Type	Immune system
Tissue	Liver tissue
Cell Type	Stellate cell
Morphology	Spindle-shaped; polygonal
Growth Properties	Adherent
Storage Conditions	For long-term cryopreservation, cryovials should be stored in liquid nitrogen at -150°C to -196°C. Storage at -80°C is restricted to short-term interim use only.

Culture Conditions and Handling

Complete Medium	Mouse Hepatic Stellate (Primary cell immortalization) Cell Complete Medium[CM-M041Y]
Population Doublings	Guaranteed to further expand for 250 population doublings
Subcultivation Ratio	First passage at 1:2 ratio, subsequent passages at 1:2 to 1:6 ratio
Dissociation Reagent	Accutase
Medium Renewal	Every 2 to 3 days
Freezing Medium	General Freezing Medium[PB180436]
Incubation Atmosphere	Air, 95%; CO ₂ , 5%
Temperature	37°C
Instructions	1. Check all containers for leakage or breakage. 2. Remove the frozen cells from the dry ice packaging and immediately transfer them to liquid nitrogen (liquid or vapor phase) for long-term cryopreservation. 3. For the first subculture, a 1:2 split ratio is recommended. Subculture at a 1:2 split ratio; cells can be subcultured for more than 10 passages. Accutase™ digestion is recommended. If trypsin digestion is used, partial cell damage and death may occur.
Subculturing Procedure	Adherent cell dissociation

Subculturing Procedure

1. Aspirate the culture medium from the T25 flask and wash the cells once with phosphate-buffered saline (PBS).
2. Add 1 mL of Accutase™ solution to the T25 flask. Gently swirl the flask to ensure the solution fully covers the bottom surface, then incubate at 37°C for 1–3 min. Examine the cells under an inverted microscope. Once cells become rounded and retracted, add 5 mL of complete growth medium to neutralize the digestion.
3. Gently pipette to mix and dissociate the cells. Subculture into a new T25 flask at the recommended split ratio, and top up with fresh complete growth medium to a final volume of 5 mL. Incubate the flask in a humidified incubator at 37°C with 5% CO₂.
4. Once the cells have fully adhered, assess cell viability and confluency under a microscope prior to downstream experiments. Subsequently, refresh the medium with fresh complete growth medium per the standard cell maintenance protocol.

Background

Mouse hepatic stellate cells (primary cell immortalization) from Procell Laboratory were isolated by lentiviral transduction of primary mouse hepatic stellate cells with the SV40T gene, followed by antibiotic selection and subculture screening. Mouse hepatic stellate cells are isolated from liver tissue. The liver is an organ primarily responsible for metabolic functions, including detoxification, glycogen storage, and synthesis of secretory proteins. It also produces bile for the digestive system. The liver is the largest internal organ of the body, located in the upper right quadrant of the abdomen, beneath the diaphragm, anterior to the right kidney, and above the stomach. It is the largest digestive gland in the body, appearing as a reddish-brown V-shaped organ. The liver is a major site of urea synthesis and plays a critical role in metabolism. Its position and morphology: the liver lies in the right upper abdomen, hidden under the diaphragm and rib cage. Most of the liver is covered by the costal margin, with only a small area exposed in the epigastric region and right costal margin, directly contacting the anterior abdominal wall. The superior surface of the liver contacts the diaphragm and the anterior abdominal wall. Hepatic stellate cells (HSCs; also known as fat-storing cells, vitamin A-storing cells, perisinusoidal cells, or Ito cells) are the primary source of extracellular matrix (ECM) in the liver. Upon activation, HSCs can further transdifferentiate into myofibroblast-like cells. Various factors that lead to liver fibrosis ultimately target HSCs. Under normal conditions, HSCs are quiescent. When the liver is injured, such as by inflammation or mechanical stress, HSCs become activated. Activated HSCs contribute to liver fibrosis and the remodeling of hepatic structure through proliferation and secretion of ECM components – a central event in liver fibrogenesis. HSCs reside in the space of Disse and are the main source of ECM synthesis and secretion, playing a pivotal role in the development of liver fibrosis. Hepatic stellate cells are liver-specific mesenchymal cells, accounting for approximately 8–13% of all liver cells. As the major ECM-producing cell population in the liver, HSCs not only secrete ECM components such as proteoglycans and glycoproteins, and synthesize a certain amount of collagenase to maintain the normal basement membrane structure, but also regulate sinusoidal microcirculation through the contraction of their processes. Furthermore, HSCs promote hepatocyte proliferation and liver regeneration by synthesizing hepatocyte growth factor (HGF), insulin-like growth factor (IGF), epidermal growth factor (EGF), and other factors.

Mouse hepatic stellate cells (primary cell immortalization) from Procell Laboratory were generated by lentiviral delivery of the SV40T gene into primary mouse hepatic stellate cells, followed by antibiotic selection and serial subculture screening. Each vial contains approximately 5×10^5 cells.

Mouse hepatic stellate cells (primary cell immortalization) isolated from Procell Laboratory exhibit positive immunofluorescence staining for α -SMA, with a purity greater than 90%. In addition, the cells test negative for HIV-1, HBV, HCV, mycoplasma, bacteria, yeast, and fungi.

Handling Recommendations for Cryopreserved Cells

Preparations Before Thawing

1. Primary cells have high nutritional demands; therefore, prepare complete culture medium in advance.
2. Preheat the complete culture medium to 37°C for 30 minutes.
3. Prepare 9 mL of complete medium within a sterile centrifuge tube to dilute the cryoprotectant present in the frozen cell suspension.
4. Use sterile gloves to protect the cryovial from contact with the water bath water, and preventing cell contamination.

Thawing Procedure

1. Retrieve the cryovial containing frozen cells from liquid nitrogen storage. Place it in sterile gloves and immediately submerge it in a 37°C water bath.
2. Thaw the cells rapidly (<1 minute) by gently swirling the vial in the 37°C water bath until only a small ice crystal remains.
3. Transfer the vial to a laminar flow hood. Wipe the exterior of the vial with 75% ethanol before opening.
4. Transfer the thawed cells dropwise into a preheated 9 mL centrifuge tube containing complete medium.
5. Centrifuge the cell suspension at approximately $250 \times g$ for 5 minutes. For cells characterized by low density and small volume, such as lymphocytes and suspended cells, the centrifugation speed can be appropriately increased to $400 \times g$ for 8 minutes.
6. After centrifugation, discard the supernatant and retain the cell pellet.
7. Gently resuspend the cells in complete growth medium, then transfer them to an appropriate culture vessel and place it in the recommended culture environment.

Notes

1. The entire recovery process should be completed as quickly as possible.
2. Select the culture vessel size based on the number of cells in the cryovial. Procell single cryovials are recommended for resuspension in 6 cm dishes or T25 flasks.
3. Minimize the time thawed cells are kept at room temperature. DMSO must be immediately diluted or removed by centrifugation.

Precautions After Receipt of Frozen Cells

1. Upon unpacking, inspect the condition of the frozen cells and dry ice, and take photographs immediately. The following after-sales service will be provided based on these photographs, including assessment of the remaining dry ice, verification that the cryovial was fully buried in dry ice, and evaluation of whether the cells thawed and refrozen during transit.
2. Upon receipt, the cells should be transferred to liquid nitrogen immediately or directly resuscitated. If liquid nitrogen is unavailable, the cells can be temporarily stored at -80°C, however, the storage period should be limited to less than one week whenever possible. Prolonged storage at -80°C may gradually reduce post-thaw cell viability, and the extent of this viability loss is unpredictable.

3. Ensure that the operator has sufficient knowledge and experience in cell culture, and the laboratory is equipped with essential instruments, including a biosafety cabinet, CO₂ incubator, inverted microscope, centrifuge, water bath. Carefully review the cell instruction manual to understand key cell characteristics including growth properties (adherent or suspension), morphology, basal medium requirements, serum concentration, cytokine supplementation, subcultivation ratio, medium renewal schedule.

4. After resuscitation, observe the cells under a microscope and record the cell status by taking photographs (1-3 images each at 100× and 200× magnification for 3 consecutive days). These images will serve as supporting documentation for follow-up services. In addition, a small aliquot of cells may be used to assess cell viability by automated cell counting or trypan blue staining.

Note: Cells should not be observed too frequently within the first 24 hours after resuscitation, as this may affect cell growth or adherence. Observation once per day is sufficient.

5. Primary cells have a limited number of passages; it is recommended that they be used as soon as possible.

Procell[®]systemProcell[®]system