Elabscience®


PE/Elab Fluor[®] 594 Anti-Mouse CD49d Antibody[R1-2]

Catalog Number: AN00422P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Mouse
Host	Rat
Isotype	Rat lgG2b, κ
Clone No.	R1-2
Isotype Control	PE/Elab Fluor [®] 594 Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09842P]
Conjugation	PE/Elab Fluor [®] 594
Conjugation Information	PE/Elab Fluor [®] 594 is designed to be excited by the blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 620 nm (e.g., a 610/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.





Staining of C57BL/6 murine splenocytes cells with PE/Elab

Fluor[®] 594 Anti-Mouse CD49d Antibody[R1-2] (filled gray histogram) or PE/Elab Fluor[®] 594 Rat IgG2b, κ Isotype Control (empty black histogram). Total viable cells were used for analysis.

Preparation & Storage	
Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	α4 integrin;VLA-4 α chain;integrin α4;ITGA4

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u> Tel: 1-832-243-6086 Email:techsupport@elabscience.com

Elabscience®

Uniprot ID Gene ID Background Q00651 16401

CD49d is a 150 kD glycoprotein, also known as α 4 integrin or VLA-4 α chain. It is a member of the integrin family, expressed on T and B cells, monocytes, eosinophils, basophils, mast cells, thymocytes, NK cells, and dendritic cells. CD49d is a heterodimer expressed with either of two β chains, β 1 (CD29) or β 7, to form the VLA-4 (integrin α 4 β 1) or LPAM-1 (integrin α 4 β 7) complexes. CD49d plays a critical role in adhesion and T cell costimulation. The primary ligands for CD49d are VCAM-1, MAdCAM-1, and fibronectin.