

A Reliable Research Partner in Life Science and Medicine

PE/Cyanine7 Anti-Human CD2 Antibody[RPA-2.10]

Catalog Number: E-AB-F1375H

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Mouse Host

Isotype Mouse IgG1, ĸ Clone No. RPA-2.10

PE/Cyanine7 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792H] Isotype Control

Conjugation

Conjugation Information PE/Cyanine7 is designed to be excited by the Blue (488 nm), Green (532 nm) and

yellow-green (561 nm) lasers and detected using an optical filter centered near 775 nm

(e.g., a 780/60 nm bandpass filter).

Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. Storage Buffer


Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



Human peripheral blood lymphocytes are stained with FITC Anti-human CD19 Antibody and PE/Cyanine7 Anti-Human CD2 Antibody[RPA-2.10] (Left). Lymphocytes are stained with FITC Anti-human CD19 Antibody and PE/Cyanine7 Mouse IgG1, κ Isotype Control (Right).

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names Erythrocytereceptor;LFA-2;LFA-3R;T-cellsurfaceantigenCD2;T11/Leu-5

Uniprot ID P06729 Gene ID 914

For Research Use Only

Fax: 1-832-243-6017 Tel: 1-832-243-6086 Toll-free: 1-888-852-8623 Email:techsupport@elabscience.com

Web:www.elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Background

CD2 is a 50 kD type I transmembrane glycoprotein also known as LFA-2, T11, and sheep red blood cell receptor (SRBC-R). This immunoglobulin superfamily member is expressed on thymocytes, T lymphocytes, NK cells, and thymic B cell subsets. The major ligand for CD2 is CD58 (also known as LFA-3). CD2 has also been reported to bind CD48, CD59, and CD15. CD2 plays a critical role in alternative T cell activation, T cell signaling, and cell-cell adhesion.

Fax: 1-832-243-6017