Recombinant Human IDH1 Protein (His Tag)

Catalog Number: PKSH032659



Note: Centrifuge before opening to ensure complete recovery of vial contents.

			•
 O C	OWI		ınn
		UU	ion

 Species
 Human

 Mol_Mass
 48.1 kDa

 Accession
 O75874

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin $< 1.0 \text{ EU per } \mu\text{g of the protein as determined by the LAL method.}$

Storage Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.

Shipping This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel


packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 μm filtered solution of 50mM Tris, 200mM NaCl, 10% glycerol,

pH8.0.

Reconstitution Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Isocitrate Dehydrogenase [NADP] Cytoplasmic (IDH1) belongs to the isocitrate and isopropylmalate dehydrogenases family. IDH1 exists as a homodimer; binding one magnesium or manganese ion per subunit. Mutations of IDH1 have been shown to cause metaphyseal chondromatosis with aciduria and are involved in the development of glioma IDH plays a role in the regeneration of NADPH for intraperoxisomal reductions; such as the conversion of 2; 4-dienoyl-CoAs to 3-enoyl-CoAs; as well as in peroxisomal reactions that consume 2-oxoglutarate; namely the α -hydroxylation of phytanic acid.

For Research Use Only