A Reliable Research Partner in Life Science and Medicine

# Recombinant Human RARRES2/TIG2 Protein (His Tag)

Catalog Number: PKSH032994

Note: Centrifuge before opening to ensure complete recovery of vial contents.

$\mathbf{T}$					
H)	es	cri	m	П	) II

Species Human

Source HEK293 Cells-derived Human RARRES2;TIG2 protein Glu21-Ser157, with an C-

terminal His

Calculated MW 16.9 kDa
Observed MW 16-20 kDa
Accession Q99969

**Bio-activity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of 20mM PB,150mM NaCl,pH7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Retinoic acid receptor responder protein 2(RARRES2) is a secreted protein that in humans is encoded by the RARRES2 gene. It is highly expressed in skin, also found in pancreas, liver, spleen, prostate, ovary, small intestine and colon. It is a chemoattractant protein that acts as a ligand for the Gprotein-coupled receptor CMKLR1. RARRES2 is secreted in an inactive form as prochemerin and is activated through cleavage of the C-terminus by inflammatory and coagulation serine proteases. It is thought to act as a cell surface receptor, found to stimulate chemotaxis of dendritic cells and macrophages to the site of inflammation. RARRES2 is inhibited in psoriatic lesions, it is activated by tazarotene in skin rafts and in the epidermis of psoriatic lesions.

### For Research Use Only