

CCL4/MIP-1 β /ACT-2, Mouse, Recombinant

Cat. No. : PCK336

General Information

Synonyms	MIP-1b: Macrophage Inflammatory Protein-1 β ;ACT-2
Species	Mouse
Expression host	E.coli
Sequence	APMGSDPPTSCCFSTSRQLHRSFVMDYYETSSLCSKPAVVFLTKRGRQICANPSEPWWTEY MSDLELN with polyhistidine tag at the N-terminus.
Accession	Q5QNV9
Tag	His-tag at the N-terminus
Mol mass	8.64 kDa
Expiration date	12 months
Bio activity	Measure by its ability to chemoattract BaF3 cells transfected with human CCR5. The ED50 for this effect is < 4 ng/mL.


Product feature

Purity	> 98% as determined by SDS-PAGE. Ni-NTA chromatography.
Endotoxin (EU/μg)	< 0.1
Storage	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
Shipping	Ice bag
Formulation	The protein was lyophilized from a 0.2 μ m filtered solution containing 1 $\times$ PBS, pH 7.4.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μ g/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Background

CCL4 act as an important pro-inflammatory cytokine during acute and chronic inflammatory responses. CCL4 also play a role as an attractant NK cells, dendritic cells, monocytes, and lymphocytes to sites of injury and inflammation. As CCL4 conducts signaling through G-protein-coupled receptor CCR5, which is a major co-receptor for M-tropic HIV strains. Thus, binding of CCL4 to CCR5 inhibits HIV entry and reduces the cell surface expression of CCR5. CCL4 has been identified as one of the major anti-HIV factors produced by CD8+ T cells.

SDS-PAGE



Proce

Procell[®]system

Procell[®]system

Procell[®]system