## **CAV1** Polyclonal Antibody

catalog number: E-AB-16228



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Synthetic peptide of human CAV1
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000
Data	
	Na 13- % %
	б— 7—
	<pre>%- %- %- %- %- Human leiomyosarcoma tissue</pre>
using CAV1 Polyclonal	<ul> <li><sup>8</sup>- <sup>n</sup>- <sup>n</sup>-</li> <li><sup>n</sup>-</li> &lt;</ul>
using CAV1 Polyclonal Calculate	<pre>%- %- %- %- %- Human leiomyosarcoma tissue</pre>
using CAV1 Polyclonal	<ul> <li><sup>8</sup>- <sup>n</sup>- <sup>n</sup>-</li> <li><sup>n</sup>-</li> &lt;</ul>
using CAV1 Polyclonal Calculate	<ul> <li><sup>8</sup>- <sup>n</sup>- <sup>n</sup>-</li> <li><sup>n</sup>-</li> &lt;</ul>

Background

The scaffolding protein encoded by this gene is the main component of the caveolae plasma membranes found in most cell types. The protein links integrin subunits to the tyrosine kinase FYN, an initiating step in coupling integrins to the Ras-ERK pathway and promoting cell cycle progression. The gene is a tumor suppressor gene candidate and a negative regulator of the Ras-p42/44 mitogen-activated kinase cascade. Caveolin 1 and caveolin 2 are located next to each other on chromosome 7 and express colocalizing proteins that form a stable hetero-oligomeric complex. Mutations in this gene have been associated with Berardinelli-Seip congenital lipodystrophy. Alternatively spliced transcripts encode alpha and beta isoforms of caveolin 1.

temperature recommended.

## For Research Use Only