A Reliable Research Partner in Life Science and Medicine

# Recombinant Human NKp30/NCR3 protein (His Tag)

Catalog Number: PDMH100415

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Species Human

**Source** HEK293 Cells-derived Human NKp30 protein Met1-Gly135, with an C-terminal His

Calculated MW 14.7 kDa
Observed MW 25 kDa
Accession 014931

**Bio-activity** Not validated for activity

### **Properties**

**Purity** > 95% as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU/mg of the protein as determined by the LAL method

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

**Reconstitution** It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Natural Cytotoxicity Triggering Receptor 3, NCR3, also known as NKp30, or CD337, is a natural cytotoxicity receptor, expressed on subsets of human peripheral blood NK cells, involved in NK cell killing of tumor cells and immature dendritic cells. The cellular ligand for NKp30 has remained elusive, but the membrane-associated heparan sulfate (HS) proteoglycans are involved in the recognition of cellular targets by NKp30 was recently reported. NKp30 is a member of the immunoglobulin superfamily and one of three existing natural cytotoxicity-triggering receptors. NKp30 is a glycosylated protein and is thought to be selectively expressed in resting and activated natural killer cells. NKp30 is a stimulatory receptor on human NK cells implicated in tumor immunity, and is capable of promoting or terminating dendritic cell maturation. NCR3 may play a role in inflammatory and infectious diseases.

### For Research Use Only