A Reliable Research Partner in Life Science and Medicine

# AF/LE Purified Anti-Human CD197 Antibody[G043H7]

catalog number: E-AB-F11590

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

**Reactivity** Human

**Immunogen** Recombinant Human CD197 protein

**Host** Mouse

**Isotype** Mouse IgG2a, κ

Clone G043H7

**Purification** >98%, Protein A/G purified


**Buffer** Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method

.

Applications	Recommended Dilution

FCM  $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$ 

#### Data



Human peripheral blood lymphocytes were stained with 0.2 μg AF/LE Purified Anti-Human CD197 Antibody[G043H7] (Right) and 0.2 μg Mouse IgG2a, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD3 FITC-conjugated Monoclonal Antibody.

## **Preparation & Storage**

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles. This preparation contains no preservatives, thus it should be handled

under aseptic conditions.

**Shipping** Ice bag

### Background

The protein encoded by this gene is a member of the Gprotein-coupled receptor family. This receptor was identified as a gene induced by the Epstein-Barr virus (EBV), and is thought to be a mediator of EBV effects on B lymphocytes. This receptor is expressed in various lymphoid tissues and activates B and T lymphocytes. It has been shown to control the migration of memory T cells to inflamed tissues, as well as stimulate dendritic cell maturation. The chemokine (C-C motif) ligand 19 (CCL19/ECL) has been reported to be a specific ligand of this receptor. Signals mediated by this receptor regulate T cell homeostasis in lymph nodes, and may also function in the activation and polarization of T cells, and in chronic inflammation pathogenesis. Alternative splicing of this gene results in multiple transcript variants.

### For Research Use Only

Fax: 1-832-243-6017