

Recombinant PRDM4/PFM1 Monoclonal Antibody

catalog number: **AN302044L**

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description



Reactivity	Human;Mouse;Rat
Immunogen	Peptide. This information is proprietary to PTMab.
Host	Rabbit
Isotype	Rabbit IgG
Clone	C121665
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications

Recommended Dilution

WB	1:1000
IHC	1:500

Data




Western Blot with PRDM4/PFM1 Monoclonal Antibody at dilution of 1:1000. Lane 1: HeLa, Lane 2: 293T, Lane 3: Mouse heart, Lane 4: Rat heart


Immunohistochemistry of paraffin-embedded Human colon using PRDM4/PFM1 Monoclonal Antibody at dilution of 1:500.

Observed-MW:100 kDa

Calculated-MW:88 kDa



Immunohistochemistry of paraffin-embedded Human pancreas using PRDM4/PFM1 Monoclonal Antibody at dilution of 1:500.



Immunohistochemistry of paraffin-embedded Human testis using PRDM4/PFM1 Monoclonal Antibody at dilution of 1:500.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2

The PRDM family members are characterized by the presence of a PR (positive regulatory) domain and multiple zinc finger (ZnFg) domains. The PR domain defines a family of transcription factors involved in cell differentiation and tumorigenesis. PRDM proteins are either epigenetic modifiers in their own right or else they recruit third party chromatin modifiers - e.g. histone deacetylases (HDACs), histone lysine MTases or histone arginine MTases - to regulate cell type-specific gene expression in various tissues (PRDM4 as an HDAC-associated transcriptional repressor that modulates cell cycle progression).