Recombinant Human TGFB1/TGF-beta 1 Protein (His Tag)

Catalog Number: PKSH033495



Description **Species** Human Mol Mass 29.3&12.8 kDa Accession P01137 **Bio-activity** Not validated for activity **Properties** > 95 % as determined by reducing SDS-PAGE. Purity < 1.0 EU per µg of the protein as determined by the LAL method. Endotoxin Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. This product is provided as lyophilized powder which is shipped with ice packs. Shipping Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Formulation Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information. Reconstitution

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| kDa | MK | R |
|-----------|----|---|
| 120 90 | | |
| 60 | | |
| 40 | | |
| 30 | | |
| 20 | - | |
| 14 | - | |

> 95 % as determined by reducing SDS-PAGE.

Background

Data

Transforming Growth Factor β -1 (TGF β -1) is a secreted protein which belongs to the TGF- β family. TGF β -1 is abundantly expressed in bone; articular cartilage and chondrocytes and is increased in osteoarthritis (OA). TGF β -1 performs many cellular functions; including the control of cell growth; cell proliferation; cell differentiation and apoptosis. The precursor is cleaved into a latency-associated peptide (LAP) and a mature TGF β -1 peptide. TGF β -1 may also form heterodimers with other TGF β family members. It has been found that TGF β -1 is frequently upregulated in tumor cells. Mutations in this gene results in Camurati-Engelmann disease.

For Research Use Only