



A Reliable Research Partner in Life Science and Medicine

# **Decorin Polyclonal Antibody(Capture/Detector)**

catalog number: AN004570P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

<b>T</b>		cript			
	20	o r		7	Int
ш	701	$\mathbf{u}$	т.	ш	LUL

**Reactivity** Human; Mouse

Immunogen Recombinant Mouse Decorin Protein expressed by Mammalian

Host Rabbit
Isotype Rabbit IgG

**Purification** Antigen Affinity Purification


**Conjugation** Unconjugated

**Buffer** Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

### Applications Recommended Dilution

ELISA Capture 2-8 μg/mL ELISA Detector 0.1-0.4 μg/mL

#### Data



Sandwich ELISA-Recombinant Mouse Decorin Protein standard curve. Background subtracted standard curve using Anti-Decorin antibody(AN004570P)(Capture), Anti-Decorin antibody(AN004570P)(Detector). The reference range value

is 31.3~2000pg/mL for mouse.

## **Preparation & Storage**

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

## Background

This gene encodes a member of the small leucine-rich proteoglycan family of proteins. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate the mature protein. This protein plays a role in collagen fibril assembly. Binding of this protein to multiple cell surface receptors mediates its role in tumor suppression, including a stimulatory effect on autophagy and inflammation and an inhibitory effect on angiogenesis and tumorigenesis. This gene and the related gene biglycan are thought to be the result of a gene duplication. Mutations in this gene are associated with congenital stromal corneal dystrophy in human patients. Decorin is a small secreted chondroitin/dermatan sulfate proteoglycan in the family of small leucine-rich proteoglycans (SLRPs). SLRP family members are characterized by N-terminal and C-terminal cysteine-rich regions which flank the central region containing 10-12 tandem leucine-rich repeats (LRR).

#### For Research Use Only

Tel: 400-999-2100 Web: www.elabscience.cn Email:techsupport@elabscience.cn