

## Recombinant Human Legumain/LGMN Protein (His Tag)

**Catalog Number:** PKSH032687

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human Legumain;LGMN protein Ile18-Tyr433, with an C-terminal His
<b>Calculated MW</b>	48.7 kDa
<b>Observed MW</b>	56 kDa
<b>Accession</b>	AAH03061.1
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Concentration</b>	Subject to label value.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
<b>Shipping</b>	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20°C.
<b>Formulation</b>	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, 10% Glycerol, pH 8.0.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Legumain is a lysosomal cysteine protease which is a member of the peptidase C13 family. Though it is found in many tissues, it is highly expressed in the kidney, heart, and placenta. Legumain has a strict specificity for hydrolysis of asparaginyl bonds and can also cleave aspartyl bonds slowly, especially under acidic conditions. Over-expression of Legumain in tumors is significant for invasion and metastasis. In addition, Legumain may be involved in the processing of proteins for MHC class II antigen presentation in the lysosomal/endosomal system and negative regulation of neuron apoptosis.

### For Research Use Only