A Reliable Research Partner in Life Science and Medicine

# Recombinant Human SCF/c-Kit Ligand Protein (His Tag)

Catalog Number: PKSH033081

Note: Centrifuge before opening to ensure complete recovery of vial contents.

		crip					
	00	CI	411	n	П	n	m
JU	$\mathbf{c}$	U		J A	ш	v.	ш

Species Human

Source HEK293 Cells-derived Human SCF; c-Kit Ligand protein Glu26-His214, with an C-

terminal His

Calculated MW22.0 kDaObserved MW30-50 kDaAccessionP21583

**Bio-activity** Measured in a cell proliferation assay using TF- 1 human erythroleukemic cells. The

 $ED_{50}$  for this effect is 2-10 ng/ml.

## **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants


before lyophilization.


Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data







Measured in a cell proliferation assay using TF- 1 human erythroleukemic cells. The ED50 for this effect is 2-10 ng/ml.

### Background

Stem Cell Factor (SCF) is a hematopoietic growth factor that exerts its activity at the early stages of hematopoiesis. SCF stimulates the proliferation of myeloid; erythroid; and lymphoid progenitors in bone marrow cultures and has been shown to act synergistically with colony stimulating factors.

### For Research Use Only