## Recombinant Human CCL27 Protein(Trx Tag)


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Catalog Number: PDEH100528



Description Species Human Source E.coli-derived Human CCL27 protein Phe25-Gly112, with an N-terminal Trx Mol Mass 29.6 kDa O9Y4X3 Accession **Bio-activity** Not validated for activity **Properties** Purity >90% as determined by reducing SDS-PAGE. Endotoxin < 10 EU/mg of the protein as determined by the LAL method Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at  $< -20^{\circ}$ C for 3 months. Shipping This product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Formulation Mannitol. Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human CCL27 proteins, 2 µg/lane of Recombinant Human CCL27 proteins was resolved with an SDS-PAGE under reducing conditions, showing bands at 29.6 KD

## Background

This gene is one of several CC cytokine genes clustered on the p-arm of chromosome 9. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The protein encoded by this gene is chemotactic for skin-associated memory T lymphocytes. This cytokine may also play a role in mediating homing of lymphocytes to cutaneous sites. It specifically binds to chemokine receptor 10 (CCR10). Studies of a similar murine protein indicate that these protein-receptor interactions have a pivotal role in T cell-mediated skin inflammation.

## For Research Use Only