

Recombinant Human Cystatin C/CST3 Protein (Human Cells, His Tag)

DIA-AN®
by Elabscience

Catalog Number:PKSH032323

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Cystatin-C;Cystatin-3;Gamma-trace;Neuroendocrine basic polypeptide;Post-gamma-globulin;CST3
Species	Human
Expression Host	HEK293 Cells
Sequence	Ser27-Ala146
Accession	P01034
Calculated Molecular Weight	14.4 kDa
Observed molecular weight	15-18 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20°C.
Formulation	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 500mM NaCl, 0.01% Tween 80, 1mM EDTA, 50% Glycerol, pH 8.0.
Reconstitution	Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Cystatin C is a member of family 2 of the cystatin superfamily. It is ubiquitous in human tissues and body fluids and mainly used as a biomarker of kidney function. Cystatin C inhibits many cysteine proteases such as papain and Cathepsins B; H; K; L and S. As an inhibitor of cysteine proteinases; Cystatin C is thought to serve an important physiological role as a local regulator of this enzyme activity. Recently; it has been studied for its role in predicting new-onset or deteriorating cardiovascular disease. It also seems to play a role in brain disorders involving amyloid (a specific type of protein deposition); such as Alzheimer's disease.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017