Recombinant Human Neuropilin-1/NRP1 (C-Avi-6His) Biotinylated

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Catalog Number: PKSH033988



Description **Species** Human Mol Mass 72.5 kDa Accession AAH07533.1 Not validated for activity **Bio-activity Properties** > 95 % as determined by reducing SDS-PAGE. Purity < 1.0 EU per µg of the protein as determined by the LAL method. Endotoxin Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. This product is provided as lyophilized powder which is shipped with ice packs. Shipping Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Formulation Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information. Reconstitution

Data			
	kDa	МК	R
	120 90	-	
		-	-
	60		
	40	-	
	20		
	30		
	20	-	

> 95 % as determined by reducing SDS-PAGE.

Background

D

Neuropilin-1 (Npn-1, previously neuropilin; also CD304) is a 130 - 140 kDa type I transmembrane (TM) glycoprotein that regulates axon guidance and angiogenesis. Two homologues, Neuropilin-1 and Neuropilin-2, are identified. Neuropilin-1 binds to semaphorin 3A, The PLGF-2 isoform of PGF, The VEGF-165 isoform of VEGF and VEGF-B. Coexpression with KDR results in increased VEGF-165 binding to KDR as well as increased chemotaxis. It may regulate VEGF-induced angiogenesis. The soluble isoform 2 binds VEGF-165 and appears to inhibit its binding to cells. NRP1 expression is regulated in EC by tumor necrosis factor-alpha, the transcription factors dHAND and Ets-1, and vascular injury. NRP1 upregulation is positively correlated with the progression of various tumors.

For Research Use Only