

Recombinant Human VDR/NR1I1 Protein (His Tag)

Catalog Number: PKSH030931

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	Baculovirus-Insect Cells-derived Human VDR/NR1I1 protein Met 1-Ser 427, with an C-terminal His
Calculated MW	50.0 kDa
Observed MW	50 kDa
Accession	P11473
Bio-activity	Not validated for activity

Properties

Purity	> 88 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 50mM Tris, 100mM NaCl, pH 8.0, 10% glycerol Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 88 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

VDR (vitamin D(1,25- dihydroxyvitamin D₃)receptor), also known as NR1I1, belongs to the NR1I family, NR1 subfamily. It is composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal ligand-binding domain. Vitamin D receptors (VDRs) are members of the NR1I family, which also includes pregnane X (PXR) and constitutive androstane (CAR) receptors, that form heterodimers with members of the retinoid X receptor family. VDRs repress expression of 1 α -hydroxylase (the proximal activator of 1,25(OH)₂D₃) and induce expression of the 1,25(OH)₂D₃ inactivating enzyme CYP24. Also, it has recently been identified as an additional bile acid receptor alongside FXR and may function to protect gut against the toxic and carcinogenic effects of these endobiotics. VDR is expressed in the intestine, thyroid and kidney and has a vital role in calcium homeostasis. It is the nuclear hormone receptor, also called transcription factor that mediates the action of vitamin D₃. Inherited mutations in the VDR gene leads to rickets.