

## PerCP/Cyanine5.5 Anti-Mouse CD8b.2 Antibody[53-5.8]

**Catalog Number:** AN00564J

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

<b>Reactivity</b>	Mouse
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG1, $\kappa$
<b>Clone No.</b>	53-5.8
<b>Isotype Control</b>	PerCP/Cyanine5.5 Rat IgG1, $\kappa$ Isotype Control[HRPN] [Product E-AB-F09822J]
<b>Conjugation</b>	PerCP/Cyanine 5.5
<b>Conjugation Information</b>	PerCP/Cyanine5.5 is designed to be excited by the blue laser (488 nm) and detected using an optical filter centered near 675 nm (e.g., a 690/50 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.


### Applications

### Recommended usage

#### FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

### Data



Staining of C57BL/6 murine splenocytes cells with FITC Anti-Mouse CD8a Antibody and PerCP/Cyanine5.5 Anti-Mouse CD8b.2 Antibody[53-5.8] (left) or PE/Cyanine7 Rat IgG1,  $\kappa$  Isotype Control (right). Total viable cells were used for analysis.

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	Lyt-3.2;Ly-3.2
<b>Uniprot ID</b>	P10300

### For Research Use Only

**Gene ID**

12526

**Background**

CD8b is the 32 kD  $\beta$  chain of CD8, also known as Lyt-3.2 or Ly-3.2. It is a member of the Ig superfamily expressed as a heterodimer with the CD8 $\alpha$  chain on a subset of MHC class I-restricted T cells and most thymocytes. CD8 is a co-receptor for the TCR complex involved in T cell activation. The antibody 53-5.8 is specific for Ly-3.2 and has low reactivity with Ly-3.1.

**For Research Use Only**