

## Recombinant Human Cathepsin Z Protein (His Tag)

**Catalog Number:** PKSH031822

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human Cathepsin Z protein Met 1-Val 303, with an C-terminal His
<b>Calculated MW</b>	33 kDa
<b>Accession</b>	Q9UBR2
<b>Bio-activity</b>	Measured by its ability to cleave the fluorogenic peptide substrate, Mca-RPPGFSAFK (Dnp)-OH, R&D Systems, Catalog # ES005. The specific activity is > 800 pmoles/min/μg.

### Properties

<b>Purity</b>	> 97 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per μg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 97 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Cathepsin Z (CTSZ); also known as Cathepsin X or CATX; belongs to the C1 family of lysosomal cysteine proteases. Its gene structure and activity properties show several unique features that distinguish it clearly from other human cysteine proteases. It has a very short pro-region that shows no similarity to those of other cathepsins and a three-residue insertion motif that forms a characteristic "mini loop". Cathepsin Z exhibits mono- and di-peptidase activity at its C-terminus; and in contrast to cathepsin B; it does not act as an endopeptidase. It is restricted to the cells of the immune system; predominantly monocytes; macrophages and dendritic cells. Cathepsin Z is widely expressed in human tissues; suggesting that this enzyme could be involved in the normal intracellular protein degradation taking place in all cell types. It is capable to cleave regulatory motifs at C-terminus affecting the function of targeted molecules. Cathepsin X may regulate also the maturation of dendritic cells; a process; which is crucial in the initiation of adaptive immunity. Furthermore; higher levels of Cathepsin Z are also found in tumour and immune cells of prostate and gastric carcinomas and in macrophages of gastric mucosa; especially after infection by *Helicobacter pylori*. Cathepsin Z is also ubiquitously distributed in cancer cell lines and in primary tumors from different sources; suggesting that this enzyme may participate in tumor progression.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017