
Recombinant Human SIRPA protein (His Tag)

Catalog Number: PDEH100822

Note: Centrifuge before opening to ensure complete recovery of vial contents.

SpeciesHumanSourceE.coli-derived Human SIRPA protein Ser105-Arg324, with an N-terminal HisCalculated MW24.1 kDaObserved MW32 kDaAccessionP78324Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	
Calculated MW24.1 kDaObserved MW32 kDaAccessionP78324Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Findotoxin< 10 EU/mg of the protein as determined by the LAL method	
Observed MW32 kDaAccessionP78324Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	
AccessionP78324Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Findotoxin< 10 EU/mg of the protein as determined by the LAL method	
Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	
Properties Purity > 95% as determined by reducing SDS-PAGE. Endotoxin < 10 EU/mg of the protein as determined by the LAL method Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -2 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquot reconstituted samples are stable at < -20°C for 3 months.	
Purity > 95% as determined by reducing SDS-PAGE. Endotoxin < 10 EU/mg of the protein as determined by the LAL method Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -2 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquot reconstituted samples are stable at < -20°C for 3 months.	
Endotoxin< 10 EU/mg of the protein as determined by the LAL methodStorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -2°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquot reconstituted samples are stable at < -20°C for 3 months.	
StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -2°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquot reconstituted samples are stable at < -20°C for 3 months.	
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquot reconstituted samples are stable at < -20 °C for 3 months.	
reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	0 to -80
1	s of
Shipping This product is provided as lyophilized powder which is shipped with ice pack	
	5.
Formulation Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%	
Mannitol.	
Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution	tion of
0.5 mg/mL. Concentration is measured by UV-Vis.	



> 95 % as determined by reducing SDS-PAGE.

Background

Data

Signal Regulatory Protein α (SIRP α) is a monomeric approximately 90 kD type I transmembrane glycoprotein. The 504 amino acid human SIRP α contains two Ig-like C1-type domains and one Ig-like V-type domain. SIRP α can express in various tissues, mainly on brain and myeloid cells, including macrophages, neutrophils, dendritic and Langerhans cells. It also can detect in neurons, smooth muscle and endothelial cells. SIRPA is an immunoglobulin-like cell surface receptor for CD47. SIRP α acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. SIRP α shows adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. SIRP α engagement generally produces a negative regulatory signal, it may mediate negative regulation of phagocytosis, mast cell activation and dendritic cell activation

For Research Use Only Toll-free: 1-888-852-8623