

Recombinant Mouse Interleukin-7/IL-7 Protein (His Tag)

Catalog Number: PKSM041098

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse Interleukin-7/IL-7 protein Glu26-Ile154 , with an C-terminal His
Calculated MW	15.9 kDa
Observed MW	22-28 kDa
Accession	P10168
Bio-activity	Measured in a cell proliferation assay using PHA-activated human peripheral blood lymphocytes (PBL). The ED ₅₀ for this effect is 60-1000 pg/ml.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

IL7, also known as interleukin 7, is a hematopoietic growth factor that belongs to the IL-7/IL-9 family. It is secreted by stromal cells in the bone marrow and thymus. IL7 stimulates the proliferation of lymphoid progenitors. It is important for proliferation during certain stages of B-cell maturation. IL7 and the hepatocyte growth factor (HGF) form a heterodimer that functions as a pre-pro-B cell growth-stimulating factor. It is found to be a cofactor for V(D)J rearrangement of the T cell receptor beta (TCRβ) during early T cell development. IL7 can be produced locally by intestinal epithelial and epithelial goblet cells and may serve as a regulatory factor for intestinal mucosal lymphocytes.

For Research Use Only