Recombinant Human CD19/Leu-12 Protein (Fc Tag)

Catalog Number: PKSH032206



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms B-Lymphocyte Antigen CD19;B-Lymphocyte Surface Antigen B4;Differentiation

Antigen CD19;T-Cell Surface Antigen Leu-12;CD19;CVID3

Species Human

Expression Host CHO Stable Cells
Sequence Pro20-Lys291

AccessionP15391Calculated Molecular Weight57.3 kDaObserved molecular weight80-95 kDaTagC-Fc

Bioactivity Immobilized Human FMC63 at 2μg/ml(100 μl/well) can bind Human CD19-Fc. The

ED₅₀ of Human CD19-Fc is 55.28 ug/ml.

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin $< 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.
 Formulation Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed man

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

CD19 is a single-pass type I membrane protein containing 2 Ig-like C2-type (immunoglobulin-like) domains. CD19 is expressed on follicular dendritic cells and B cells. In fact; it is present on B cells from earliest recognizable B-lineage cells during development to B-cell blasts but is lost on maturation to plasma cells. CD19 primarily acts as a B cell co-receptor in conjunction with CD21 and CD81. Upon activation; the cytoplasmic tail of CD19 becomes phosphorylated; which leads to binding by Src-family kinases and recruitment of PI-3 kinase. CD19 Assembles with the antigen receptor of B

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

Recombinant Human CD19/Leu-12 Protein (Fc Tag)

Catalog Number: PKSH032206



lymphocytes in order to decrease the threshold for antigen receptor-dependent stimulation. Defects in CD19 are the cause of immunodeficiency common variable type 3 (CVID3) which is a primary immunodeficiency characterized by antibody deficiency; hypogammaglobulinemia; recurrent bacterial infections and an inability to mount an antibody response to antigen.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com