# Acetyl-P53 (Lys381) Polyclonal Antibody

catalog number: E-AB-20287



Note: Centrifuge before opening to ensure complete recovery of vial contents.

		crip			
- 1	00	CPI	m	17	٦m
J			174	, T. U	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,

Reactivity Human; Mouse; Rat

**Immunogen** Synthesized peptide derived from human p53 around the acetylation site of K381

Host Rabbit Isotype IgG


Purification Affinity purification
Conjugation Unconjugated

**buffer** Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein

protectant and 50% glycerol.

Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:100-1:300
IF	1:200-1:1000

#### Data



Western Blot analysis of HepG2, 293T-UV, Hela, AD293T,

Hela-UV cells using p53 (Acetyl Lys381) Polyclonal

Antibody at dilution of 1:1000.

Observed-MV:53 kDa Calculated-MV:44 kDa

# **Preparation & Storage**

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack,upon receipt, store it immediately at the

temperature recommended.

### Background

The protein encoded by this gene is a cytoplasmic kinase that is structurally similar to the yeast Ste20p kinase, which acts upstream of the stress-induced mitogen-activated protein kinase cascade. The encoded protein can phosphorylate myelin basic protein and undergoes autophosphorylation. A caspase-cleaved fragment of the encoded protein has been shown to be capable of phosphorylating histone H2B. The particular phosphorylation catalyzed by this protein has been correlated with apoptosis, and it's possible that this protein induces the chromatin condensation observed in this process.

## For Research Use Only