



A Reliable Research Partner in Life Science and Medicine

Mouse GM-CSF Antibody Pair Set

Catalog No. E-KAB-0322 Applications ELISA

Synonyms GMCSF, CSF2

Kit components & Storage

Title	Specifications	Storage
Mouse GM-CSF Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Mouse GM-CSF Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0322)	
		Mouse GM-CSF Capture Antibody	Mouse GM-CSF Detection Antibody (Biotin)
Immunogen	Immunogen	Recombinant Mouse GM-CSF	Recombinant Mouse GM-CSF protein
Information		protein	
	Swissprot	P01587	
Product details	Reactivity	Mouse	Mouse
	Host	Rat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Mouse GM-CSF in ELISAs.	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com





A Reliable Research Partner in Life Science and Medicine

Applications

Mouse GM-CSF Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4μg/mL	Mouse GM-CSF Capture Antibody	
Capture			Aist
ELISA Detection	1:1000-1:10000	Mouse GM-CSF Detection Antibody (Biotin)	O.01 1000 10000 Mouse GM-CSF concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

CSF2 (Colony Stimulating Factor 2) is a Protein Coding gene. Diseases associated with CSF2 include Mucositis and Candida Glabrata. Among its related pathways are RET signaling and IL-15 Signaling Pathways and their Primary Biological Effects in Different Immune Cell Types. GO annotations related to this gene include cytokine activity and granulocyte macrophage colony-stimulating factor receptor binding. The protein encoded by this gene is a cytokine that controls the production, differentiation, and function of granulocytes and macrophages. The active form of the protein is found extracellularly as a homodimer. This gene has been localized to a cluster of related genes at chromosome region 5q31, which is known to be associated with interstitial deletions in the 5q- syndrome and acute myelogenous leukemia. Other genes in the cluster include those encoding interleukins 4, 5, and 13.