

## NME1/NDKA Monoclonal Antibody

**catalog number:** AN200029N

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Reactivity</b>	Human
<b>Immunogen</b>	Recombinant Human NME1 / NDKA protein
<b>Host</b>	Mouse
<b>Isotype</b>	IgG1
<b>Clone</b>	6A9
<b>Purification</b>	Protein A
<b>Buffer</b>	0.2 µm filtered solution in 20 mM MES, 100 mM NaCl, 10% Trehalose, 0.02% Tween 80, pH 6.0


### Applications

Applications	Recommended Dilution
ICC/IF	1:20-1:100
FCM	1:25-1:100

### Data



Flow cytometric analysis of Human NME1 expression on HeLa cells. The cells were stained with purified anti-Human NME1, then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells.



Immunofluorescence analysis of Human NME1 in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with Mouse anti-Human NME1 Monoclonal Antibody (1:60) at 37°C 1 hour. Then cells were stained with the Alexa Fluor® 594-conjugated Goat Anti-mouse IgG secondary antibody (red) and counterstained with DAPI for nuclear staining (blue). Positive staining was localized to cytoplasm.

### Preparation & Storage

<b>Storage</b>	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
<b>Shipping</b>	Ice bag

### Background

### For Research Use Only

This gene (NME1) was identified because of its reduced mRNA transcript levels in highly metastatic cells. Nucleoside diphosphate kinase (NDK) exists as a hexamer composed of 'A' (encoded by this gene) and 'B' (encoded by NME2) isoforms. Mutations in this gene have been identified in aggressive neuroblastomas. Two transcript variants encoding different isoforms have been found for this gene. Co-transcription of this gene and the neighboring downstream gene (NME2) generates naturally-occurring transcripts (NME1-NME1), which encodes a fusion protein comprised of sequence sharing identity with each individual gene product.