

# Recombinant Human CA9 Protein( Fc Tag)

Catalog Number: PDMH100227

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Species</b>	Human
<b>Source</b>	Mammalian-derived Human CA9 protein Gln38-Asp414,with an C-terminal Fc
<b>Mol_Mass</b>	66.3 kDa
<b>Accession</b>	Q16790
<b>Bio-activity</b>	Not validated for activity

## Properties

<b>Purity</b>	> 90% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

## Data



SDS-PAGE analysis of Human CA9 proteins, 2 µg/lane of Recombinant Human CA9 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 66.3 KD

## Background

Carbonic anhydrases IX (CA IX), also known as membrane antigen MN or CA9, is a member of the carbonic anhydrase (CA) family and may be involved in cell proliferation and cellular transformation. CAs are zinc metalloenzymes that catalyze the reversible hydration of carbon dioxide ( $H_2O + CO_2 = H^+ + HCO_3^-$ ) and thus participate in a variety of biological and physical processes. CA IX protein is expressed primarily in carcinoma cells lines, and the expression is cell density dependent and has been shown to be strongly induced by hypoxia, accordingly facilitates adaptation of tumor cells to hypoxic conditions. It is involved in tumorigenesis through many pathways, such as pH regulation and cell adhesion control. CA IX is used as a marker of tumor hypoxia and as a new therapeutic target for many human carcinomas and cancers.

## For Research Use Only