A Reliable Research Partner in Life Science and Medicine

Recombinant Human HGF Protein (His Tag)

Catalog Number: PKSH032538

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source HEK293 Cells-derived Human HGF protein Gln32-Ser728, with an C-terminal His

 Mol_Mass
 26&53.7 kDa

 Accession
 P14210

Bio-activity Measured by its ability to induce IL-11 secretion by Saos- 2 human osteosarcoma

cells. The ED50 for this effect is 0.3-1. 5 ng/ml.

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 0.01 EU per μg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

 Shipping
 This product is provided as lyophilized powder which is shipped with ice packs.

 Formulation
 Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 500mM NaCl, pH 8.0.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Hepatocyte growth factor/scatter factor (HGF/SF) is a paracrine cellular growth, motility and morphogenic factor. It belongs to the peptidase S1 family and Plasminogen subfamily, contains 4 kringle domains, 1 PAN domain and 1 peptidase S1 domain. HGF regulates cell growth, cell motility, and morphogenesis by activating a tyrosine kinase signaling cascade after binding to the proto-oncogenic c-Met receptor. HGF is secreted by mesenchymal cells and acts as a multi-functional cytokine on cells of mainly epithelial origin. Its ability to stimulate mitogenesis, cell motility, and matrix invasion gives it a central role in angiogenesis, tumorogenesis, and tissue regeneration.

For Research Use Only

Fax: 1-832-243-6017