## **Elabscience**®

## Recombinant Human Bruton Tyrosine Kinase/BTK Kinase Protein (His Tag)

## Catalog Number: PKSH030407

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	Baculovirus-Insect Cells-derived Human Bruton Tyrosine Kinase/BTK Kinase protein
	Met 1-Ser 659, with an C-terminal His
Calculated MW	77.8 kDa
Accession	NP_000052.1
<b>Bio-activity</b>	The specific activity was determined to be 115 nmol/min/mg using Poly(Glu, Tyr)4:1
	peptide as substrate.
Properties	
Purity	> 85 % as determined by reducing SDS-PAGE.
Concentration	Subject to label value.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at $<$ -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel
	packs. Upon receipt, store it immediately at $< -20^{\circ}$ C.
Formulation	Supplied as sterile solution of 20mM Tris, 500mM NaCl, 10% glycerol, pH 7.0
Data	
	KDa MK R
	116
	66.2
	45.0
	35.0
	25.0
	18.4

> 85 % as determined by reducing SDS-PAGE.

14.4 \_\_\_\_\_

## Background

Bruton's tyrosine kinase (or BTK) is a type of kinase protein expressed in B lymphocytes and T cells. BTK contains a PH domain which binds phosphatidylinositol(3,4,5)-trisphosphate (PIP3). After binding to PIP3, BTK is induced to phosphorylate phospholipase C, which in turn hydrolyzes PIP2 into two second messagers, IP3 and DAG, which then modulate the activity of downstream proteins during B-cell signaling. Btk is also found implicated in the primary immunodeficiency disease X-linked agammaglobulinemia(Bruton's agammaglobulinemia). BTK played a key role in B-cell maturation as well as mast cell activation through the high-affinity IgE receptor. Patients with X-linked agammaglobulinemia have normal pre-B cell populations in their bone marrow but these B-cells can not mature and enter the circulation.