

Recombinant Human FAM3B (C-Fc)

Catalog Number: PKSH033848

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

| | |
|----------------------|--|
| Species | Human |
| Source | HEK293 Cells-derived Human FAM3B protein Glu30-Ser235, with an C-terminal Fc |
| Calculated MW | 50.0 kDa |
| Observed MW | 50-65 kDa |
| Accession | P58499 |
| Bio-activity | Not validated for activity |

Properties

| | |
|-----------------------|---|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. |
| Reconstitution | Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information. |

Data



Background

FAM3B, also known as Pancreatic-derived factor (PANDER), is an islet-specific secreted cytokine specifically expressed at high levels in the islets of Langerhans of the endocrine pancreas. FAM3B can induce apoptosis of alpha and beta cells in a dose- and time-dependent manner. Previous studies showed that FAM3B regulates glucose and lipid metabolism through interaction with liver and endocrine pancreas. FAM3B silencing activates both extrinsic and intrinsic apoptotic pathways. In general, silencing FAM3B promoted p53 phosphorylation and induced p53 accumulation by decreasing Mdm2 expression, which resulted in apoptotic cell death.

For Research Use Only