Elabscience®

Mouse CX3CL1 Antibody Pair Set

Catalog No.E-KAB-0326ApplicationsELISASynonymsNeurotactin, ABCD-3, C3Xkine, CXC3, CXC3C, NTN, NTT, SCYD1

Kit components & Storage

Title	Specifications	Storage
Mouse CX3CL1 Capture Antibody	1 vial, 100 µ g	Store at -20° C for one year.
		Avoid freeze / thaw cycles.
Mouse CX3CL1 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0326)	
		Mouse CX3CL1 Capture Antibody	Mouse CX3CL1 Detection Antibody (Biotin)
Immunogen	Immunogen	Recombinant Mouse CX3CL1	Recombinant Mouse CX3CL1 protein
Information		protein	
	Swissprot	O35188	
Product details	Reactivity	Mouse	Mouse
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Mouse CX3CL1 in ELISAs.	

For Research Use Only

Elabscience®

Applications

Mouse CX3CL1 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4µg/mL	Mouse CX3CL1 Capture Antibody	
Capture			
ELISA	1:1000-1:10000	Mouse CX3CL1 Detection	Optical Density
Detection		Antibody (Biotin)	
			Mouse CX3CL1 concentration(ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

This gene encodes a large cytokine protein of 373 amino acids, it contains multiple domains and is the only known member of the CX3C chemokine family. It is also commonly known under the names fractalkine (in humans) and neurotactin (in mice). The polypeptide structure of CXC3L1 differs from the typical structure of other chemokines. CX3CL1 is produced as a long protein (with 373-amino acid in humans) with an extended mucin-like stalk and a chemokine domain on top. The mucin-like stalk permits it to bind to the surface of certain cells. However a soluble (90 kD) version of this chemokine has also been observed. Soluble CX3CL1 potently chemoattracts T cells and monocytes, while the cell-bound chemokine promotes strong adhesion of leukocytes to activated endothelial cells, where it is primarily expressed. CX3CL1 elicits its adhesive and migratory functions by interacting with the chemokine receptor CX3CR1. Its gene is located on human chromosome 16 along with some CC chemokines known as CCL17 and CCL22.

For Research Use Only