
## Recombinant Mouse CD6/TP120 Protein (His Tag)

## Catalog Number: PKSM041351

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Mouse
Source	HEK293 Cells-derived Mouse CD6/TP120 protein Leu18-Gly396, with an C-terminal
	His
Calculated MW	41.9 kDa
Observed MW	55-95 kDa
Accession	Q91WN5
Bio-activity	Not validated for activity
Properties	
Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}C$ for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	



## Background

## **Elabscience**®

CD6 is a member of the group B scavenger receptor cysteinerich(SRCR) superfamily. CD6 is a type I membrane glycoprotein and contains three extracellularSRCR domains. CD6 is expressed at low levels on immature thymocytes and at high levels on mature thymocytes. The majority of peripheral blood T cells, a subsetof B cells, and a subset of neuronal cells express CD6. Mouse CD6 is a 626 amino acid (aa) protein with a 24 aa sequence, a 372 aa extracellular domain, and a204 aa cytoplasmic region. The role of CD6 has not been fully elucidated. However, it appears to play a role as both a costimulatorymolecule in T cell activation and as an adhesion receptor.CD6/ALCAM interactions have been postulated to play a role in thymocyte development. The CD6 intracellular domain contains regions that can interact with SH2 or SH3 containing proteins. However, the signaling pathways have not been elucidated.