

Recombinant Mouse CXCL1/Lymphotoxin1/I-TAC Protein(Gst Tag)

Catalog Number: PDEM100187

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	E.coli-derived Mouse CXCL1/Lymphotoxin1/I-TAC protein Phe22-Met100, with an N-terminal Gst
Calculated MW	34.5 kDa
Observed MW	40 kDa
Accession	Q9JHH5
Bio-activity	Not validated for activity

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Mouse CXCL1/Lymphotoxin1/I-TAC proteins, 2µg/lane of ReCombinant Mouse CXCL1/Lymphotoxin1/I-TAC proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 40 KD

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.7

I-TAC, also known as CXCL1/Lymphotactin1/I-TAC, is a small cytokine belonging to the CXC chemokine family. It is highly expressed in peripheral blood leukocytes, pancreas and liver, with moderate levels in thymus, spleen and lung and low expression levels were in small intestine, placenta and prostate. The I-TAC chemokine elicits its effects on its target cells by interacting with the cell surface chemokine receptor CXCR3, with a higher affinity than do the other ligands for this receptor, CXCL9 and CXCL1/Lymphotactin0. I-TAC is chemotactic for activated T cells. The CXCL1/Lymphotactin1/I-TAC gene is located on human chromosome 4 along with many other members of the CXC chemokine family.