

CD5/Cluster of Differentiation 5 Monoclonal Antibody

catalog number: **AN200254P**

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Rat
Immunogen	Recombinant Rat CD5/Cluster of Differentiation 5 Protein
Host	Mouse
Isotype	IgG1
Clone	12B6
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS


Applications Recommended Dilution

ICC/IF	1:20-1:100
FCM	1:25-1:100

Data



Flow cytometric analysis of Rat CD5 expression on SD rat splenocytes. Cells were stained with purified anti-Rat CD5, then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells.



Immunofluorescence analysis of Rat CD5 in rat spleen cells. Cells were fixed with 4% PFA, blocked with 10% serum, and incubated with mouse anti-ratCD5 monoclonal antibody (dilution ratio 1:60) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-mouse IgG secondary antibody (green). Positive staining was localized to Cell membrane.

Preparation & Storage

Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag

Background

For Research Use Only

T-cell differentiation antigen CD6, also known as TP12 and CD6, is a single-pass type I membrane protein which contains three SRCR domains. CD6/TP12 is a cell surface glycoprotein expressed primarily on T cells, it may function as a costimulatory molecule and may play a role in autoreactive immune responses. CD6/TP12 is expressed by thymocytes, mature T-cells, a subset of B-cells known as B-1 cells, and by some cells in the brain. CD6 ligand termed CD166 (previously known as activated leukocyte cell adhesion molecule, ALCAM) has been identified and shown to be expressed on activated T cells, B cells, thymic epithelium, keratinocytes, and in rheumatoid arthritis synovial tissue. CD6/TP12 binds to activated leukocyte cell adhesion molecule (CD166), and is considered as a costimulatory molecule involved in lymphocyte activation and thymocyte development. CD6/TP12 partially associates with the TCR/CD3 complex and colocalizes with it at the center of the mature immunological synapse (IS) on T lymphocytes. During thymic development CD6-dependent signals may contribute both to thymocyte survival, and to the overall functional avidity of selection in both man and mouse.