Purified Anti-Human CD80 Antibody[2F4]

catalog number: AN003450P



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human CD80 protein

Host Mouse

Isotype Mouse IgGl, κ

Clone 2F4

Purification >98%, Protein A/G purified


Conjugation Unconjugated

buffer PBS, pH 7.2. Contains 0.05% proclin 300.

Applications Recommended Dilution

FCM $2 \mu g/mL(1\times10^5-5\times10^5 \text{ cells})$

Data



Raji were stained with 0.2 μg Purified Anti-Human CD80 Antibody[2F4] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-

Mouse IgG Secondary Antibody.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

Background

B7-1 and B7-2, together with their receptors CD28 and CTLA-4, constitute one of the dominant co-stimulatory pathways that regulate T- and B-cell responses. Although both CTLA-4 and CD28 can bind to the same ligands, CTLA-4 binds to B7-1 and B7-2 with a 20-100 fold higher affinity than CD28 and is involved in the down-regulation of the immune response. B7-1 is expressed on activated B cells, activated T cells, and macrophages.

For Research Use Only