A Reliable Research Partner in Life Science and Medicine

# **UBE2D1 Polyclonal Antibody**

catalog number: E-AB-60521

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human; Mouse; Rat

Immunogen Recombinant fusion protein of human UBE2D1 (NP 003329.1).

Host Rabbit Isotype IgG


Purification Affinity purification


Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

#### **Recommended Dilution Applications**

WB 1:500-1:2000 IHC 1:50-1:200

#### Data







UBE2D1 Polyclonal Antibody at dilution of 1:1000.

Western blot analysis of extracts of various cell lines using Immunohistochemistry of paraffin-embedded Rat brain using UBE2D1 Polyclonal Antibody at dilution of 1:100 (40x lens).

## Observed-MV:14 kDa Calculated-MV:16 kDa





Immunohistochemistry of paraffin-embedded Rat spleen using UBE2D1 Polyclonal Antibody at dilution of 1:100 (40x lens).

Immunohistochemistry of paraffin-embedded Mouse spleen using UBE2D1 Polyclonal Antibody at dilution of 1:100 (40x lens).

#### Preparation & Storage

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Storage

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

### **Background**

#### For Research Use Only

Rev. V1.6

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

Elabscience®

The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. This enzyme is closely related to a stimulator of iron transport (SFT), and is upregulated in hereditary hemochromatosis. It also functions in the ubiquitination of the tumor-suppressor protein p53 and the hypoxia-inducible transcription factor HIF1alpha by interacting with the E1 ubiquitin-activating enzyme and the E3 ubiquitin-protein ligases. Two transcript variants encoding different isoforms have been found for this gene.

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Tel: 1-832-243-6086 Email:techsupport@elabscience.com