

A Reliable Research Partner in Life Science and Medicine

## Recombinant Clusterin/Apolipoprotein J/Apo-J Monoclonal Antibody

catalog number: AN300045P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human

Immunogen Recombinant Human Clusterin / Apolipoprotein J / Apo-J protein


Host Rabbit Isotype lgG Clone A1155 **Purification** Protein A

Buffer 0.2 µm filtered solution in PBS


#### **Applications Recommended Dilution**

1:20-1:100 ICC/IF 1:25-1:100 **FCM** 

#### Data



were stained with purified anti-Human CLU, then a FITCconjugated second step antibody. The histogram were derived from gated events with the forward and side lightscatter characteristics of intact cells.



Flow cytometric analysis of Human CLU on A549 cells. Cells Immunofluorescence analysis of Human CLU in Hela cells. Cells were fixed with 4% PFA, permeabilzed with 1% Triton X-100 in PBS, blocked with 10% serum, and incubated with Rabbit anti-Human CLU Monoclonal Antibody (1:60) at 37°C 1 hour. Then cells were stained with the Alexa Fluor® 488conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to cytoplasm.

### **Preparation & Storage**

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

**Shipping** Ice bag

**Background** 

Toll-free: 1-888-852-8623 Fax: 1-832-243-6017 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com Rev. V1.2

# Elabscience®

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

Clusterin, also known as complement-associated protein SP-40, Complement cytolysis inhibitor, Apolipoprotein J, Testosterone-repressed prostate message 2, Aging-associated gene 4 protein, CLU and APOJ, is a secreted protein which belongs to the clusterin family. Clusterin/Apolipoprotein J/Apo-J is an enigmatic glycoprotein with a nearly ubiquitous tissue distribution and an apparent involvement in biological processes ranging from mammary gland involution to neurodegeneration in Alzheimer's disease. Its major form, a heterodimer, is secreted and present in physiological fluids, but truncated forms targeted to the nucleus have also been identified. Clusterin/Apolipoprotein J/ Apo-J is a widely distributed glycoprotein with a wide range of biologic properties. A prominent and defining feature of clusterin is its marked induction in such disease states as glomerulonephritis, cystic renal disease, renal tubular injury, neurodegenerative conditions, atherosclerosis, and myocardial infarction. Upregulation of clusterin mRNA and protein levels detected in diverse disease states and in in vitro systems have led to suggestions that it functions in membrane lipid recycling, in apoptotic cell death, and as a stress-induced secreted chaperone protein, amongst others.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.2