Recombinant Human NFASC/Neurofascin protein (His Tag)

Catalog Number: PKSH030521

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	HEK293 Cells-derived Human NFASC/Neurofascin protein Met 1-Gln939, with an C-
	terminal His
Calculated MW	104.4 kDa
Observed MW	114-119 kDa
Accession	O94856-12
Bio-activity	Not validated for activity
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $<$ -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	
	KDa MK R
	116
	66.2
	45.0
	35.0
	25.0
	25.0

> 95 % as determined by reducing SDS-PAGE.

18.4 14.4

Background

NFASC, also known as neurofascin, belongs to the immunoglobulin superfamily, L1/neurofascin/NgCAM family. It contains 5 fibronectin type-III domains and 6 Ig-like C2-type (immunoglobulin-like) domains. NFASC functions in neurite outgrowth, neurite fasciculation, and organization of the axon initial segment (AIS) and nodes of Ranvier on axons during early development. Both the AIS and nodes of Ranvier contain high densities of voltage-gated Na+ (Nav) channels which are clustered by interactions with cytoskeletal and scaffolding proteins including this protein, gliomedin, ankyrin 3 (ankyrin-G), and betaIV spectrin. NFASC links the AIS extracellular matrix to the intracellular cytoskeleton.

For Research Use Only Toll-free: 1-888-852-8623