

Human MMP-7 Antibody Pair Set

Catalog No.	E-KAB-0501	Applications	ELISA
Synonyms	MMP7;MPSL1;PUMP-1;Matrilysin		

Kit components & Storage

Title	Specifications	Storage
Human MMP-7 Capture Antibody	1 vial, 100 µg	Store at -20℃ for one year. Avoid freeze/thaw cycles.
Human MMP-7 Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20℃ for one year. Avoid freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Product Information

Items		Characteristic (E-KAB-0501)	
		Human MMP-7 Capture Antibody	Human MMP-7 Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Human MMP-7 protien	Recombinant Human MMP-7 protien
	Swissprot	P09237	
Product details	Reactivity	Human	Human
	Host	Mouse	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50% glycerol; pH 7.5	PBS with 0.04% Proclin 300; 1% protective protein; 50% glycerol; pH 7.5
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Human MMP-7 in ELISAs.	

For Research Use Only

Applications

Human MMP-7 Sandwich ELISA Assay

	Recommended Concentration/Dilution	Reagent	Images
ELISA Capture	0.5-4 µg/mL	Human MMP-7 Capture Antibody	
ELISA Detection	1:1000-1:10000	Human MMP-7 Detection Antibody (Biotin)	

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The enzyme encoded by this gene degrades proteoglycans, fibronectin, elastin and casein and differs from most MMP family members in that it lacks a conserved C-terminal protein domain. The enzyme is involved in wound healing, and studies in mice suggest that it regulates the activity of defensins in intestinal mucosa. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3.

For Research Use Only