Elabscience®


Elab Fluor[®] Violet 450 Anti-Human CD4 Antibody[RPA-T4]

Catalog Number: E-AB-F1109Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Host	Mouse
lsotype	Mouse IgG1, κ
Clone No.	RPA-T4
Isotype Control	Elab Fluor [®] Violet 450 Mouse IgG1, к Isotype Control[MOPC-21] [Product E-AB- F09792Q]
Conjugation	Elab Fluor [®] Violet 450
Conjugation Information	Elab Fluor [®] Violet 450 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Human peripheral blood lymphocytes are stained with APC

Anti-Human CD19 Antibody and Elab Fluor[®] Violet 450 Anti-Human CD4 Antibody (Left). Lymphocytes are stained with

APC Anti-Human CD19 Antibody and Elab Fluor[®] Violet 450 Mouse IgG1, κ Isotype Control (Right).

Preparation & Storage		
Storage	Keep as concentrated solution.	
	This product can be stored at 2-8°C for 12 months. Please protect exposure to light and do not freeze.	cted from prolonged
Shipping	Ice bag	
Antigen Information		
Alternate Names	CD4;T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein C	CD4
Uniprot ID	P01730	
For Research Use Only	/	
Toll-free: 1-888-852-8623	Tel: 1-832-243-6086	Fax: 1-832-243-6017
Web:www.elabscience.com	Email:techsupport@elabscience.com	Rev. V1.6

Elabscience®

HHV-7).

Gene ID	920
Background	CD4, also known as T4/Leu-3, is a 55 kD single-chain type I transmembrane
	glycoprotein and member of the immunoglobulin superfamily. It is expressed on most
	thymocytes, helper T cells, type II NKT cells, and monocytes/macrophages. CD4 is part

of the TCR/CD3 complex, binds to β 2 domain from the MHC class II molecule, and participates in TCR signal transduction. CD4 is the receptor of IL-16 and is a

coreceptor for the human immunodeficiency virus (HIV) and human herpes virus 7 (

For Research Use Only