

Recombinant Human FUT8 Protein (aa 68-575, His Tag)

Catalog Number: PKSH031148

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	Baculovirus-Insect Cells-derived Human FUT8 protein Arg 68-Lys 575, with an C-terminal His
Mol_Mass	60.0 kDa
Accession	Q9BYC5-1
Bio-activity	Measured by its ability to hydrolyze the donor substrate GDP fucose. The specific activity is > 0.75 pmoles/min/μg.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 8.0, 10% glycerol Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Alpha (1,6) fucosyltransferase 8, also known as FUT8, is a member of the glycosyltransferase family.

Fucosyltransferases are the enzymes transferring fucose from GDP-Fuc to Gal in an alpha1,2-linkage and to GlcNAc in alpha1,3-linkage, alpha1,4-linkage, or alpha1,6-linkage. All fucosyltransferases utilize the same nucleotide sugar, their specificity reside in the recognition of the acceptor and in the type of linkage formed. Fucosyltransferases share some common structural and catalytic features. On the basis of protein sequence similarities, these enzymes can be classified into four distinct families: (1) the alpha-2-fucosyltransferases, (2) the alpha-3-fucosyltransferases, (3) the mammalian alpha-6-fucosyltransferases, and (4) the bacterial alpha-6-fucosyltransferases. The alpha-3-fucosyltransferases constitute a distinct family as they lack the consensus peptide, but some regions display similarities with the alpha-2 and alpha-6-fucosyltransferases.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017