

## Recombinant Human PKC iota/PRKCI Protein (GST Tag)

**Catalog Number:** PKSH030411

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	Baculovirus-Insect Cells-derived Human PKC iota/PRKCI protein Met 10-Val 596, with an C-terminal GST
<b>Calculated MW</b>	93.5 kDa
<b>Observed MW</b>	100 kDa
<b>Accession</b>	NP_002731.4
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 88 % as determined by reducing SDS-PAGE.
<b>Concentration</b>	Subject to label value.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
<b>Shipping</b>	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
<b>Formulation</b>	Supplied as sterile solution of 50mM Tris, 100mM NaCl, 0.5mM GSH, 0.5mM PMSF, 10% glycerol, pH 7.4

### Data



> 88 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Protein kinase C iota type, also known as Atypical protein kinase C-lambda/iota, aPKC-lambda/iota and PRKCI, is a cytoplasm, membrane and nucleus protein which belongs to the protein kinase superfamily, AGC Ser/Thr protein kinase family and PKC subfamily. PRKCI contains one AGC-kinase C-terminal domain, one OPR domain, one phorbol-ester/DA G-type zinc finger and one protein kinase domain. PRKCI is predominantly expressed in lung and brain, but also expressed at lower levels in many tissues including pancreatic islets. It is highly expressed in non-small cell lung cancers. PRKCI is a calcium-independent, phospholipid-dependent, serine- and threonine-specific kinase. It may play a role in the secretory response to nutrients. PRKCI is involved in cell polarization processes and the formation of epithelial tight junctions. It is implicated in the activation of several signaling pathways including Ras, c-Src and NF-kappa-B pathways. PRKCI functions in both pro- and anti-apoptotic pathways. It functions in the RAC1/ERK signaling required for transformed growth. PRKCI plays a role in microtubule dynamics through interaction with RAB2A and GAPDH and recruitment to vesicular tubular clusters (VTCs). PRKCI might be a target for novel lipid activators that are elevated during nutrient-stimulated insulin secretion.