

A Reliable Research Partner in Life Science and Medicine

AF/LE Purified Anti-Human CD1a Antibody[OKT-6]

catalog number: E-AB-F11260

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

| Reactivity | Human |
|------------|-------|
| Host | Mouse |

Isotype Mouse IgG1, κ

Clone OKT-6

Conjugation None (AF/LE)


Buffer Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL

method.

Applications Recommended Dilution

FCM $2 \mu g/mL(0.5 \times 10^6 - 1 \times 10^6 \text{ cells})$

Data



Molt-4 were stained with 0.2 μg AF/LE Purified Anti-Human CD1a Antibody[OKT-6] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by FITC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

Storage Storage, avoid freeze /

thaw cycles. This preparation contains no preservatives, thus it should be handled

under aseptic conditions.

Shipping Ice bag

Background

CD1a is a 49 kD member of the immunoglobulin superfamily also known as T6 and R4. It is a type I membrane glycoprotein with structural similarities to MHC class I and is non-covalently associated with β2-microglobulin. CD1a plays a role in non-peptide glycolipid antigen presentation to CD1-restricted T cells. It is expressed on cortical double positive and single positive thymocytes, Langerhans cells, and dendritic cells. In addition to antigen presentation, CD1a has been implicated in thymic T cell development.