

NFκB-p65 Monoclonal Antibody

catalog number: E-AB-22016

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Human;Mouse
Immunogen	Synthetic Peptide
Host	Mouse
Isotype	IgG
Clone	9B4
Purification	Protein A purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein protectant and 50% glycerol.


Applications

WB

Recommended Dilution

1:1000-3000

Data



Western Blot analysis of 1) Hela, 2) 293T, 3) HepG2 cells using NFκB-p65 Monoclonal Antibody at dilution of 1:2000.

Observed-MW:65 kDa

Calculated-MW:60 kDa

Preparation & Storage

Storage

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping

The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

Proteins encoded by the v-Rel viral oncogene and its cellular homolog, c-Rel, are members of a family of transcription factors that include the two subunits of the transcription factor NFκB (p50 and p65) and the Drosophila maternal morphogen, dorsal. Both proteins specifically bind to DNA sequences that are the same or slight variations of the 10 bp κB sequence in the immunoglobulin κ light chain enhancer. This same sequence is also present in a number of other cellular and viral enhancers. The DNA binding activity of NFκB is activated and NFκB is subsequently transported from the cytoplasm to the nucleus in cells exposed to mitogens or growth factors. cDNAs encoding precursors for two distinct proteins of the same size have been described, designated p105 and p100. The p105 precursor contains p50 at its N-terminus and a C-terminal region that when expressed as a separate molecule, designated pDI, binds to p50 and regulates its activity.

For Research Use Only