

Purified Anti-Mouse CD172a Antibody[P84], Functional Grade

catalog number: E-AB-F12860

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Immunogen	Recombinant Mouse CD172a protein
Host	Rat
Isotype	Rat IgG1, κ
Clone	P84
Purification	>98%, Protein A/G purified
Buffer	Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method.

Applications


Recommended Dilution

FCM	2 $\mu\text{g}/\text{mL}$ (0.5×10^6 - 1×10^6 cells)
Block	Reported in the literature

Data



C57/BL6 Mouse bone marrow were stained with 0.2 μg Purified Anti-Mouse CD172a Antibody[P84], Functional Grade (Right) and 0.2 μg Rat IgG1, κ Isotype Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Rat IgG Secondary Antibody.



Monomer purity $\geq 95\%$ as determined by analytical size-exclusion chromatography (SEC)

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.
Shipping	Ice bag

Background

For Research Use Only

Immunoglobulin-like cell surface receptor for CD47. Acts as a docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. Supports adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. May play a key role in intracellular signaling during synaptogenesis and in synaptic function. Involved in the negative regulation of receptor tyrosine kinase-coupled cellular responses induced by cell adhesion, growth factors or insulin. Mediates negative regulation of phagocytosis, mast cell activation and dendritic cell activation. CD47 binding prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells. Plays a role in antiviral immunity and limits new world arenavirus infection by decreasing virus internalization. Receptor for THBS1. Interaction with THBS1 stimulates phosphorylation of SIRPA. In response to THBS1, involved in ROS signaling in non-phagocytic cells, stimulating NADPH oxidase-derived ROS production.

None (Azide-Free, Low Endotoxin) are perfectly suited to be used in culture or in vivo (for nonhuman studies) for functional assays blocking, neutralizing, activation or depletion where the presence of azide may damage cells or exogenous endotoxin may signal or activate cells.

Application References

Tadahiko Yanagita, et al. JCI Insight. 2017 Jan 12;2(1):e89140.