Arginase 1/ARG1 Polyclonal Antibody(Capture/Detector)

catalog number: AN000310P



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-					
- 10	AC	cri	m	TO	m
\mathbf{L}		CII		LLU.	

Reactivity Mouse

Immunogen Recombinant Mouse Arginase 1/ARGl protein expressed by E.coli

Host Rabbit
Isotype Rabbit IgG

Purification Antigen Affinity Purification


Conjugation Unconjugated

buffer Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

Applications Recommended Dilution

ELISA Capture 2-8 μg/mL **ELISA Detector** 0.1-0.4 μg/mL

Data



Sandwich ELISA-Recombinant Mouse Arginase 1/ARG1 protein standard curve.Background subtracted standard curve using Arginase 1/ARG1 antibody(AN000310P)

(Capture),Arginase 1/ARG1 antibody(AN000310P)

(Detector). in sandwich ELISA. The reference range value for Recombinant Mouse Arginase 1/ARG1 protein is 78.13-5000

pg/mL.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

Arginase is the focal enzyme of the urea cycle hydrolysing l-arginine to urea and l-ornithine. Emerging studies have identified arginase in the vasculature and have implicated this enzyme in the regulation of nitric oxide (NO) synthesis and the development of vascular disease. Arginase also redirects the metabolism of l-arginine to l-ornithine and the formation of polyamines and l-proline, which are essential for smooth muscle cell growth and collagen synthesis. Arginase is encoded by two recently discovered genes (Arginase I and Arginase II). In most mammals, Arginase 1 (ARGI) also known as Arginase, liver, which functions in the urea cycle, and is located primarily in the cytoplasm of the liver. The second isozyme, Arginase II, has been implicated in the regulation of the arginine/ornithine concentrations in the cell. It is located in mitochondria of several tissues in the body, with most abundance in the kidney and prostate. It may be found at lower levels in macrophages, lactating mammary glands, and brain.

For Research Use Only