



A Reliable Research Partner in Life Science and Medicine

Rat VEGFR-2 Antibody Pair SetSet

Catalog No. E-KAB-0391 Applications ELISA

Synonyms VEGFR2, CD309, FLK1

Kit components & Storage

Title	Specifications	Storage
Rat VEGFR-2 Capture Antibody	1 vial, 100 μ g	Store at -20°C. Avoid freeze/thaw
		cycles.
Rat VEGFR-2 Detection Antibody	1 vial, 50 μL	Store at -20°C. Avoid freeze/thaw
(Biotin)		cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0391)	
		Rat VEGFR-2 Capture Antibody	Rat VEGFR-2 Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Rat VEGFR-2 protein	Recombinant Rat VEGFR-2 protein
Information	Swissprot	O08775	
Product details	Reactivity	Rat	Rat
	Host	Rabbit	Rabbit
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300; 1%
		glycerol, pH 7.4	protective protein; 50% glycerol; pH
			7.4
	Purify	Protein A	Protein A & Antigen Affinity
	Specificity	Detects Rat VEGFR-2 in ELISAs.	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com



A Reliable Research Partner in Life Science and Medicine

Applications

Rat VEGFR-2 Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Rat VEGFR-2 Capture Antibody	
Capture			10
			Optical Donalty
ELISA	1:1000-1:10000	Rat VEGFR-2 Detection Antibody	ğ. 0 0.1
Detection		(Biotin)	
			0.01 0.1 1 10 100 Rat VEGFR-2 Concentration (ng/mL)
			Kat VEGEN-2 Concentration (ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Vascular endothelial growth factor (VEGF) is a major growth factor for endothelial cells. This gene encodes one of the two receptors of the VEGF. This receptor, known as kinase insert domain receptor, is a type III receptor tyrosine kinase. It functions as the main mediator of VEGF-induced endothelial proliferation, survival, migration, tubular morphogenesis and sprouting. The signalling and trafficking of this receptor are regulated by multiple factors, including Rab GTPase, P2Y purine nucleotide receptor, integrin alphaVbeta3, T-cell protein tyrosine phosphatase, etc.. Mutations of this gene are implicated in infantile capillary hemangiomas.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com