LDHA Polyclonal Antibody

catalog number: E-AB-19937



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen Synthetic peptide of human LDHA

Host Rabbit Isotype IgG


Purification Antigen affinity purification

Conjugation Unconjugated

buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:30-1:150

Data




Western blot analysis of A375 and Hela cell lysates using LDHA Polyclonal Antibody at dilution of 1:400

Immunohistochemistry of paraffin-embedded Human breast cancer tissue using LDHA Polyclonal Antibody at dilution of $1:40(\times 200)$

Observed-MV:Refer to figures

Calculated-MV:37 kDa



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using LDHA Polyclonal Antibody at dilution of $1:40(\times 200)$

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

LDHA Polyclonal Antibody

catalog number: E-AB-19937



LDHA, also named as LDH-M and NY-REN-59, is an enzyme which catalyzes the inter-conversion of pyruvate and L-lactate with concomitant inter-conversion of NADH and NAD+. LDHA is found in most somatic tissues, though predominantly in muscle tissue and tumours, and belongs to the lactate dehydrogenase family. It has long been known that many human cancers have higher LDHA levels compared to normal tissues. It has also been shown that LDHA plays an important role in the development, invasion and metastasis of malignancies. Mutations in LDHA have been linked to exertional myoglobinuria. LDHA has some isoforms with MW 26-40 kDa.