

Recombinant Mouse 2B4/CD244 Protein (His Tag)

Catalog Number: PKSM041150

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse 2B4/CD244 protein Gln20-Asn221, with an C-terminal His
Calculated MW	23.5 kDa
Observed MW	35-60 kDa
Accession	Q07763
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Reconstitution	Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information.

Data



Background

For Research Use Only

Natural killer cell receptor 2B4 (2B4/CD244) is a 66 kDa type I transmembrane glycoprotein in the SLAM subgroup of the CD2 protein family. SLAM family proteins have an extracellular domain (ECD) with two or four Ig-like domains and at least two cytoplasmic immunoreceptor tyrosine-based switch motifs (ITSMs). 2B4 interacts with CD48, while other SLAM family proteins interact in a homophilic manner. The mouse 2B4 cDNA encodes a 397 amino acid (aa) precursor that includes a 19 aa signal sequence, a 207 aa ECD with one Ig-like V-type and one C2-type Ig-like domain, a 21 aa transmembrane segment, and a 150 aa cytoplasmic domain with four ITSMs. Within the ECD, mouse 2B4 shares 46% and 68% aa sequence identity with human and rat 2B4, respectively. 2B4/CD48 signaling cooperates with other receptor systems to either promote or inhibit NK and CD8+ T cell activation. The inhibitory activities are distinct from those of MHC I restricted inhibitory NK cell receptors. Ligation of 2B4 with antibodies or CD48 constructs can directly trigger inhibitory signaling or disrupt an inhibitory interaction, leading to cellular activation. 2B4 can also induce signaling through CD48.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017