

A Reliable Research Partner in Life Science and Medicine

PE Anti-Mouse CD2 Antibody[RM2-5]

Catalog Number: E-AB-F1387D

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

IsotypeRat lgG2b, λ Clone No.RM2-5

Isotype Control PE Rat IgG2b, λ Isotype Control[G013B8] [Product AN00565D]

Conjugation PE

Conjugation Information PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green

(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42

nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein


protectant.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 murine splenocytes cells with PE Anti-Mouse CD2 Antibody[RM2-5] (filled gray histogram) or PE Rat IgG2b,λ Isotype Control (empty black histogram). Total viable cells were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names LFA-2;T11;Ly-37;SRBC-R

Uniprot ID P08920

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Gene ID

Background

12481

CD2 is a 45-58 kD type I transmembrane glycoprotein, also known as LFA-2, T11 or Ly-37. It is a member of the Ig superfamily. Mouse CD2 is primarily expressed on T cells, B cells, thymocytes and NK cells. It is a ligand for CD48 and is involved in T cell activation and differentiation.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com